

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Successful pharmacogenetics-based optimization of unboosted atazanavir plasma exposure in HIV-positive patients: a randomized, controlled, pilot study (the REYAGEN study)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1522605> since 2016-05-31T13:50:52Z

Published version:

DOI:10.1093/jac/dkv208

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is a pre-copyedited, author-produced PDF of an article accepted for publication in Journal of Antimicrobial Chemotherapy following peer review. The version of record “J Antimicrob Chemother. 2015 Jul 14. doi:10.1093/jac/dkv208 [Epub ahead of print]” is available online at: <http://jac.oxfordjournals.org/content/early/2015/07/13/jac.dkv208.long>

33 **Successful Pharmacogenetics-based Optimization of Unboosted Atazanavir Plasma Exposure**
34 **in HIV-positive Patients: a Randomized, Controlled, Pilot Study (The REYAGEN Study).**

35

36 °Bonora S¹, °Rusconi S², *Calcagno A¹, Bracchi M^{1,3}, Viganò O², Cusato J¹, Lanzafame M⁴,
37 Trentalange A¹, Marinaro L¹, Siccardi M⁵, D'Avolio A¹, Galli M², and Di Perri G¹.

38

39 °These two authors equally contributed to the study

40

41 ¹*Unit of Infectious Diseases, Department of Medical Sciences, University of Torino, Torino, Italy;*

42 ²*Department of Infectious Diseases, Ospedale Luigi Sacco, University of Milano, Milano, Italy;* ³*St.*

43 *Stephen's Centre, Chelsea and Westminster Hospital, London, United Kingdom;* ⁴*Unit of Diagnosis*

44 *and Therapy of HIV Infection, 'G.B.Rossi' Hospital, 37134 Verona, Italy;* ⁵*Department of*

45 *Pharmacology, University Of Liverpool, Liverpool, United Kingdom.*

46

47

48 **Eudract number:** 2009-014216-35

49 **Running Head:** PG-based adjustment of atazanavir

50

51 **Type of article:** Original article

52

53 **Key words:** pharmacogenetics; pharmacokinetics; HIV; atazanavir; ABCB1; PXR; SLCO1B1.

54

55 **Word count:** 1492 (abstract 151)

56

57 **Figures:** 1 **Tables:** 1

58 **Supplementary material:** 2 Tables, 2 Figures, CONSORT checklist.

59

60

61 ***Corresponding Author:**

62 Andrea Calcagno
63 Unit of Infectious Diseases
64 Department of Medical Sciences
65 University of Torino
66 Amedeo di Savoia Hospital
67 C.so Svizzera 164
68 10149 Torino, Italy
69 +390114393884
70 +390114393942
71 andrea.calcagno@unito.it
72

73

74

75

76

77

78

79

80

81

82

83

84

85

86 **Synopsis**

87 **Background:** Atazanavir without ritonavir, despite efficacy and tolerability, shows low plasma
88 concentrations that warrant optimization.

89 **Methods:** In a randomized, controlled, pilot trial, stable HIV-positive patients on
90 atazanavir/ritonavir (with tenofovir/emtricitabine) were switched to atazanavir. In the standard dose
91 arm atazanavir was administered as 400 mg once-daily, while according to patients' genetics (*PXR*,
92 *ABCB1* and *SLCO1B1*) in the pharmacogenetic arm: patients with unfavourable genotypes received
93 atazanavir 200 mg twice-daily.

94 **Results:** Eighty patients were enrolled with balanced baseline characteristics. Average atazanavir
95 exposure was 253 ng/mL (150-542) in the pharmacogenetic arm versus 111 ng/mL (64-190) in the
96 standard arm ($p<0.001$); 28 patients in the pharmacogenetic arm (75.7%) had atazanavir exposure
97 >150 ng/mL versus 14 patients (38.9%) in the standard arm ($p=0.001$). Immunovirological and
98 laboratory parameters had a favourable outcome throughout the study with non-significant
99 differences between study arms.

100 **Conclusions:** Atazanavir plasma exposure is higher when the schedule is chosen according to the
101 patient's genetic profile.

102

103

104

105

106

107

108 INTRODUCTION

109 In the lifelong perspective of anti-HIV treatment, individual tailoring of the antiretroviral regimen is
110 going to be increasingly required. Although never formally approved in Europe, the use of
111 atazanavir without concurrent intake of ritonavir has been shown to be effective and well tolerated
112 in two induction-maintenance clinical trials of relevant size and several retrospective studies.¹⁻⁴
113 However in a significant proportion of patients the pharmacokinetic (PK) exposure of atazanavir
114 might be potentially insufficient to guarantee long-term HIV inhibition.^{5,6} atazanavir lower
115 exposure when combined with tenofovir disoproxil fumarate has been shown in healthy volunteers
116 but subsequently found to be less relevant in HIV-positive patients.⁷⁻⁹ atazanavir pharmacokinetics
117 is significantly influenced by genetic polymorphisms in the region coding for the pregnane X
118 receptor (*PXR*, controlling the expression of several genes involved in drug metabolism and
119 transport); additionally polymorphisms in *ABCB1* (coding for P-glycoprotein) and *SLCO1B1*
120 (coding for OATP1B1) were shown to have a comparable effect on atazanavir exposure.¹⁰⁻¹²
121 Furthermore we observed that the pharmacokinetic exposure of atazanavir was significantly
122 improved when administered 200 mg twice-daily instead of 400 mg once-daily.¹³

123 We report here the results of a randomized comparative study on the clinical use of unboosted
124 atazanavir with or without pharmacogenetic guide in patients also taking co-formulated
125 tenofovir/emtricitabine.

126

127 METHODS

128 HIV-positive adult patients on treatment with atazanavir/ritonavir (300/100 mg) plus
129 tenofovir/emtricitabine with HIV RNA <50 copies/mL for at least six months were eligible for
130 enrolment at two sites in Italy. Switch to atazanavir was proposed for toxicity/tolerability or for
131 simplification, according to physicians' evaluation in clinical practice. Exclusion criteria were:

132 previous virological failure, genotypic resistance-associated mutations, ongoing opportunistic
133 infections/neoplasias, liver cirrhosis, chronic renal failure, self-reported adherence <90% (Visual
134 Scale) and consumption of potentially interacting drugs.

135 The study was approved by the institutional review board at both participating centres, and each
136 participant provided signed informed consent before enrolment; the procedures were in accordance
137 with the ethical standards of the Helsinki Declaration of 1975 (as revised in 1983).

138 The study was a randomized, controlled, open-label, pilot trial. Patients were randomized 1:1
139 (block randomization) to either standard-dose arm ["SD"; atazanavir 400 mg *once daily*] or
140 pharmacogenetic-based arm ["PG"; atazanavir 400 mg *once daily* in patients with favourable
141 genetic profile or atazanavir 200 mg *twice daily* in patients with unfavourable genetic profile]. At
142 enrolment genomic DNA was extracted using QIAamp whole blood mini kit (Qiagen, Valencia,
143 CA, USA) and genotyping was conducted by real time-based allelic discrimination with the use of
144 standard methods (BIORAD, Milano, Italy). The following single nucleotide polymorphisms were
145 analysed: C63396T in *PXR* (rs2472677), C3435T in *ABCB1* (rs1045642) and C521T in *SLCO1B1*
146 (rs4149056). *PXR* 63396 TT, *ABCB1* 3435 CT/TT and *SLCO1B1* 521 TT were codified as 1
147 (associated with lower plasma concentrations). On the basis of the PG results patients were given a
148 score (min zero - max three) and a different dosing schedule according to favourable (≤ 1) or
149 unfavourable genetic profiles (≥ 2).

150 Primary end point was the prevalence of atazanavir average trough concentrations (geometric mean
151 of the first three determinations at weeks 4, 8 and 12) above 150 ng/mL (suggested target plasma
152 level) in the two arms. Secondary end points were the comparison of the proportion of patients with
153 HIV RNA <50 copies/mL and of the changes in indirect bilirubin, total cholesterol, LDL-
154 cholesterol, HDL-cholesterol and triglycerides at 48 weeks.

155 atazanavir trough plasma concentrations [12/24 hours after drug intake according to drug schedule
156 (\pm two hours)] were measured by a previously validated HPLC-PDA (Photo Diode Array) method
157 and performed in Torino.¹⁴

158 A sample size of 80 patients (40 per group) was calculated to provide a statistical power of at least
159 80%, in order to identify a difference in mean atazanavir Ctrough below the MEC of 150 ng/mL
160 between the two study arms. It was assumed a 20% of atazanavir Ctrough under MEC in the PG
161 arm, and a 50% in the control arm from previous studies results.¹⁰⁻¹² Standard non-parametric tests
162 were used for all analysis and performed using SPSS 20.0 software for Mac (SPSS, IBM Inc.).

163

164 **RESULTS**

165 Eighty patients were enrolled (2009-2011): demographic and immunovirological characteristics
166 were well balanced between study arms (Table 1). Patients' disposition is shown in Figure S1: no
167 subject dropped out of the study due to toxicity, virological failure or major clinical events. The
168 prevalence of single nucleotide polymorphisms is reported in Table 1; all variants were in Hardy-
169 Weinberg equilibrium. 27 patients in the PG arm received atazanavir 200 mg twice daily.

170 Atazanavir plasma trough concentrations are shown in Figure S2 and Table S1. Atazanavir Ctrough
171 was slightly higher at baseline in the PG arm [1034 ng/mL (592-1935) versus. 587 ng/mL (77-
172 1290), Mann-Whitney $p=0.06$] as compared to SD arm; it was significantly higher at every time
173 point after randomization ($p<0.001$ for all comparisons, Mann-Whitney) in the PG arm.

174 Geometric mean of week 4 to 12 atazanavir Ctroughs was 253 ng/mL (150-542) in the PG arm
175 versus 111 ng/mL (64-190) in the SD arm, favouring the former ($p<0.001$, Mann-Whitney). As for
176 the primary endpoint 28 patients in the PG arm (75.7%) had an average atazanavir Ctrough above
177 150 ng/mL versus 14 patients (38.9%) in the SD arm ($p=0.001$, RR 4.89, 95%CI 1.79-13.38) (Fig.
178 1).

179 No difference in plasma HIV-RNA <50 copies/mL was observed in 37 patients (100%) in the PG
180 arm versus 33 patients (97%) in the SD arm at week 48. Three patients (8.1%) and 4 patients
181 (11.7%) in the PG and SD arm presented a viral blip during the study (p=0.703, Fisher's exact test).
182 Patients in both arms had similar CD4+ T lymphocytes recovery at week 48: 39 cells/mm³ in the
183 PG versus 53 cells/mm³ in the SD arm (p=0.744, Mann-Whitney).

184 At 48 weeks significant decreases (all p<0.05, Wilcoxon's) in safety markers were noted as
185 compared to baseline: no significant differences between study arms were found (Mann-Whitney),
186 (Table S2).

187

188 **DISCUSSION**

189 In this pilot, randomized and controlled study we found that the pharmacokinetic exposure of
190 atazanavir, when co-administered with tenofovir/emtricitabine was significantly higher and closer
191 to the desired target concentration when the frequency of administration was chosen according to
192 the patient's genetic profile. The proportion of patients with atazanavir Ctrough above the cut-off
193 concentration rose from 40% (previous studies and the standard arm) to 75.7% (study arm) when
194 the frequency of atazanavir administration (400 mg once daily or 200 mg twice daily) was decided
195 on the basis of the individual genotypic profile.¹⁰⁻¹² Although not all patients had a Ctrough level
196 above the pre-specified cut-off value of 150 ng/mL, the pharmacokinetic exposure in the study arm
197 was found significantly more appropriate than in patients in the control arm. In the PG arm baseline
198 atazanavir levels were higher than those recorded in the SD arm: it is possibly due to unbalanced
199 factors between study arms (such as *CYP3A5* genotype and adherence levels) and unexpected
200 atazanavir exposures according to genotype (Supp.Tab.1) may support this hypothesis.¹⁵ It must
201 however be considered that the 150 ng/mL threshold resulted from the analysis of a moderately
202 experienced population of HIV-infected patients that was no longer formally re-assessed in

203 treatment-naïve patients: it appears possible that it could be lower in patients not harbouring virus
204 with resistance associated mutations and after achieving viral suppression.^{6,16,17} The documented
205 higher intracellular accumulation of atazanavir as compared to other PIs might also support this
206 hypothesis.^{18,19} No significant difference in the prevalence of viral control or in the changes in
207 safety markers between study arms was seen: it is possible the longer follow-up may be required to
208 observe the effect of improved pharmacokinetic exposure or that lower atazanavir concentrations
209 may be adequate.

210 Independently of study arm atazanavir-based regimens were well tolerated and associated with
211 improved safety profiles. Even if the drug is nowadays less used given the availability of safe and
212 very compact antiretroviral regimens it may be very useful in the long-term treatment of HIV-
213 positive patients. The absence of ritonavir (associated with side effect even at low doses) and the
214 uncommon incidence of hyperbilirubinemia (being the main determinant of atazanavir/ritonavir
215 inferior performance in naïve patients) support the attractiveness of atazanavir-containing
216 regimens.²⁰ Even if the need for genetic testing prior to start atazanavir might not be commonly
217 accepted it can be a tool for avoiding unnecessary treatment interruptions and side effects.²¹
218 Although some patients (those with unfavourable genetic profile) would necessitate to take the drug
219 twice daily instead of once daily, the advantage in terms of side effects reduction might compensate
220 the higher frequency of administration.

221 We have to recognize some limitations of this study: the limited sample size, the restricted number
222 of included genetic polymorphisms as well as a casual impaired factors distribution between the
223 study arms, the potential need for therapeutic drug monitoring even in the PG-based arm.

224 Once in a lifetime performed genetic testing offers the possibility to know in advance the likelihood
225 of an individual patient to achieve a more appropriate atazanavir pharmacokinetic exposure and to

226 choose the frequency of administration accordingly; if confirmed, this observation supports the use
227 of pharmacogenetics for treatment tailoring in atazanavir-receiving HIV-positive patients.

228

229 **Acknowledgements**

230 This work was presented at the 20th Conference on Retroviruses and Opportunistic Infections, 3-6
231 March 2013, Atlanta, GE, USA. P556 and at the 14th Conference of the European AIDS Clinical
232 Society, Bruxelles, Belgium. 16-19 October 2013.

233

234 **Funding**

235 This study was funded by an unrestricted grant from Bristol Meyers-Squibb, the manufacturer of
236 atazanavir. The funding source had no role in data collection, analysis, interpretation, trial design,
237 patient recruitment, writing of the manuscript or the decision to submit it for publication. The
238 corresponding author had full access to all the data in the study and had final responsibility for the
239 decision to submit for publication.

240 **Transparency declarations**

241 **S. B.** has received grants, travel grants, and consultancy fees from Abbott, Boehringer-Ingelheim,
242 BMS, Gilead Sciences, GlaxoSmithKline (GSK), MSD, Pfizer, and Janssen-Cilag. **S.R.** has
243 received grants, travel grants and speaker's honoraria from Abbott, Bristol-Myers Squibb (BMS),
244 Merck Sharp & Dohme (MSD), and Janssen-Cilag. **A. C.** has received grants, travel grants and
245 speaker's honoraria from Abbott, Bristol-Myers Squibb (BMS), Merck Sharp & Dohme (MSD),
246 and Janssen-Cilag. **M.G.** has received grants, travel grants and speaker's honoraria from Abbott,
247 Bristol-Myers Squibb (BMS), Merck Sharp & Dohme (MSD), and Janssen-Cilag. **G. D. P.** has
248 received grants, travel grants, and consultancy fees from Abbott, Boehringer-Ingelheim, BMS,

249 Gilead Sciences, GSK, MSD, Pfizer, Roche, and Tibotec (Johnson & Johnson). All other authors
250 report no potential conflicts.

251 SB, SR, MS, AD and GDP designed the study and contributed to data collection. AC performed
252 data interpretation and statistical analysis and generated the random allocation sequence. SR, MB,
253 OV, ML, AT, LM contributed to data collection. AC and MB drafted the first version of the
254 manuscript and finalized the manuscript. JC, MS and AD performed the pharmacokinetic and
255 pharmacogenetic analysis and revised the technical details of the paper. SB, SR, GDP and MG
256 contributed to study design, supervision and critical revision of the manuscript for intellectual
257 content. All authors read and approved the final manuscript.

258

259 **References**

- 260 1. Squires KE, Young B, DeJesus E, et al. ARIES 144 week results: durable virologic
261 suppression in HIV-infected patients simplified to unboosted
262 atazanavir/abacavir/lamivudine. *HIV Clin Trials*. 2012;**13**(5):233-44.
- 263 2. Ghosn J, Carosi G, Moreno S, et al. Unboosted atazanavir-based therapy maintains control
264 of HIV type-1 replication as effectively as a ritonavir-boosted regimen. *Antivir Ther*.
265 2010;**15**(7):993-1002.
- 266 3. Pavie J, Porcher R, Torti C et al. Efficacy and safety of a switch to unboosted atazanavir in
267 combination with nucleoside analogues in HIV-1-infected patients with virological
268 suppression under antiretroviral therapy. *J Antimicrob Chemother* 2011;**66**(10): 2372-8.
- 269 4. Bonora S, Calcagno A, Vigano O et al. Efficacy, Tolerability and Virological Consequences
270 of Long-Term Use of Unboosted Atazanavir Plus 2 NRTIs in HIV-Infected Patients. *Curr*
271 *HIV Res*. 2014;**12**(5):339-46.

5. Goutelle S, Baudry T, Gagnieu MC, et al. Pharmacokinetic-pharmacodynamic modeling of unboosted Atazanavir in a cohort of stable HIV-infected patients. *Antimicrob Agents Chemother.* 2013;**57**(1):517-23.
6. Rekić D, Röshammar D, Bergstrand M, et al. External validation of the bilirubin-atazanavir nomogram for assessment of atazanavir plasma exposure in HIV-1-infected patients. *AAPS J.* 2013;**15**(2):308-15.
7. Taburet AM, Piketty C, Chazallon C, et al. Interactions between atazanavir-ritonavir and tenofovir in heavily pretreated human immunodeficiency virus-infected patients. *Antimicrob Agents Chemother.* 2004;**48**(6):2091-6.
8. Calcagno A, Bonora S, Tettoni MC, et al. Tenofovir coadministration is not associated with lower unboosted atazanavir plasma exposure in the clinical setting. *J Acquir Immune Defic Syndr.* 2009;**52**(3):431-2.
9. Fournier C, Higgins N, Thomas R, et al. Negligible effect of tenofovir on atazanavir trough concentrations and genotypic inhibitory quotients in the presence and absence of ritonavir. *Ther Drug Monit.* 2013;**35**(2):264-9.
10. Siccardi M, D'Avolio A, Baietto L, et al. Association of a single-nucleotide polymorphism in the pregnane X receptor (PXR 63396C-->T) with reduced concentrations of unboosted atazanavir. *Clin Infect Dis.* 2008;**47**(9):1222-5.
11. Schipani A, Siccardi M, D'Avolio A, et al. Population pharmacokinetic modeling of the association between 63396C->T pregnane X receptor polymorphism and unboosted atazanavir clearance. *Antimicrob Agents Chemother.* 2010;**54**(12):5242-50.
12. Siccardi M, D'Avolio A, Bonora S, et al. Combined effect of SLCO1B1 521T>C, PXR 63396C>T, and ABCB1 3435C>T on the achievement of therapeutic concentrations of unboosted Atazanavir. In: *Abstracts of the 10th International Workshop on Clinical Pharmacology of HIV Therapy, Amsterdam, The Netherlands, April 2009.* Abstract O-04.

13. Bonora S, Gonzalez de Requena D, D'Avolio A, et al. Pharmacokinetics of switching unboosted atazanavir coadministered with tenofovir disoproxil fumarate from 400 mg once daily to 200 mg twice daily in HIV-positive patients. *Antivir Ther*. 2011;**16**(4):499-504.
14. D'Avolio A, Baietto L, Siccardi M, et al. An HPLC-PDA method for the simultaneous quantification of the HIV integrase inhibitor raltegravir, the new nonnucleoside reverse transcriptase inhibitor etravirine, and 11 other antiretroviral agents in the plasma of HIV-infected patients. *Ther Drug Monit*. 2008; **30**(6):662-9.
15. Savic RM, Barrail-Tran A, Duval X, et al. Effect of Adherence as Measured by MEMS, Ritonavir Boosting, and CYP3A5 Genotype on Atazanavir Pharmacokinetics in Treatment-Naive HIV-Infected Patients. *Clin Pharmacol and Ther*. 2012; **30**(6):662-9.
16. Zhu L, Liao S, Child M, et al. Pharmacokinetics and inhibitory quotient of atazanavir/ritonavir versus lopinavir/ritonavir in HIV-infected, treatment-naive patients who participated in the CASTLE Study. *J Antimicrob Chemother*. 2012;**67**(2):465-8.
17. Lescure FX, Poirier JM, Meynard JL, et al. Factors predictive of virological failure on atazanavir in 310 HIV-infected patients. *AIDS*. 2010;**24**(10):1593-5.
18. Zehnacker L, Abe E, Mathez D, Alvarez JC, Leibowitch J, Azoulay S. Plasma and Intracellular Antiretroviral Concentrations in HIV-Infected Patients under Short Cycles of Antiretroviral Therapy. *AIDS Res Treat*. 2014;**2014**:724958.
19. D'Avolio A, Carcieri C, Cusato J, et al. Intracellular accumulation of atazanavir/ritonavir according to plasma concentrations and OATP1B1, ABCB1 and PXR genetic polymorphisms. *J Antimicrob Chemother*. 2014;**69**(11):3061-6.
20. Lennox JL, Landovitz RJ, Ribaud HJ, et al. Efficacy and tolerability of 3 nonnucleoside reverse transcriptase inhibitor-sparing antiretroviral regimens for treatment-naive volunteers infected with HIV-1: a randomized, controlled equivalence trial. *Ann Intern Med*. 2014;**161**(7):461-71.

21. Nettles RE, Child MJ, Bertz RJ, Schnittman S. Gilbert syndrome and the development of
antiretroviral therapy-associated hyperbilirubinemia: genetic screening is unnecessary. J
Infect Dis. 2006;**193**(11):1611-2

Characteristic	Standard arm (n= 40)	dose Pharmacogenetic arm (n= 40)	p values
Age (years): median (IQR)	43 (37-47)	44 (38-50)	0.424
Male gender: n (%)	28 (70%)	30 (75%)	0.783
Ethnicity: n (%)			
White	37 (92.5%)	34 (85%)	0.487
Black	1 (2.5%)	3 (7.5%)	
Other	2 (5%)	3 (7.5%)	
BMI (Kg/m ²): median (IQR)	22.9 (20.2-25.3)	23.9 (21-26.2)	0.421
Duration of HIV infection (years): median (IQR)	5.9 (3.7-12.4)	7.3 (3.7-12.3)	0.665
CD4+ T lymphocytes (cells/mm ³): median (IQR)	541 (428-628)	467 (320-600)	0.063
CD4+/CD8+ T lymphocytes ratio: median (IQR)	0.65 (0.53-1.1)	0.60 (0.5-1.29)	0.864
Hepatitis B surface antigen positive: n (%)	6 (15%)	1 (2.5%)	0.049
Hepatitis C antibody positive: n (%)	8 (20%)	8 (20%)	0.823
Single nucleotide polymorphisms: n (%)			
<i>PXR 63396</i> TT	12 (30%)	10 (25%)	0.848
<i>ABCB1 3435</i> CT/TT	28 (70%)	29 (72.5%)	0.364
<i>SLCO1B1 521</i> TT	30 (75%)	33 (82.5%)	0.848
Favorable pharmacogenotypic score (<=1): n (%)	14 (35%)	13 (32.5%)	0.797

348

349 **Table 1. Demographics, immunovirological and pharmacogenetic characteristics of**
350 **randomized patients.** Values were compared between the two arms using Chi-square (Fisher's
351 exact test where appropriate) for categorical values and Mann-Whitney test for continuous variable;
352 two-sided p values are shown in the last column. "IQR": interquartile range.

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370 **Figures:**

371


372

373

374

375

376



377

378 **Figure 1. Atazanavir average concentration (weeks 4 to 12) according to study arm.** Symbols
 379 indicate geometric mean of trough concentration obtained at weeks 4, 8 and 12; the horizontal lines
 380 represent median values. The gray boxes represent the percentage of patients with average exposure
 381 above 150 ng/mL.

382